

**Skills Competency Severe Allergic Reaction (Anaphylaxis) Medication Administration
Epinephrine Auto-Injectors 9/1/2021**

Annual skill verification is recommended by a registered nurse, medical provider or a skilled and willing parent.

Student's name: _____ Grade/Teacher: _____

Person trained: _____ Position: _____ Initials: _____

Person training: _____ Position: _____ Initials: _____

Skills:	Initial Demonstration		Return Demonstration				
	Date	Date	Date	Date	Date	Date	Date
1. Identify that symptoms of a severe allergic reaction are present and that based on the child's severe allergic reaction emergency plan, medication needs to be given.							
2. Call for assistance. Ask that another school staff personnel call 911 or emergency medical services and notify parent. Remove other students from the area, as necessary.							
3. Gather needed supplies and place on a clean surface near the student.							
4. Quickly talk to the student and evaluate the situation. Prepare the student.							
5. Position student, providing as much privacy as possible. The injector can be given through clothing in most circumstances.							
6. Quickly review the medication administration form, the medical provider or district protocol/plan form, and the parent/guardian consent form, if available, as you review the "5 Rights" of medication administration.							

7. Check the epinephrine autoinjector and order to be sure it is:							
a. For the right child							
b. The right medication							
c. The right dose, an approximation of the child's weight may be needed, if using a stock epinephrine auto-injector							
d. Being given at the right time and							
e. Being given by the right route.							
8. Be sure to check the medication to ensure that it is not expired							
9. Wash your hands, if possible. In a true emergency situation, there may not be time to do this step.							
10. Put on gloves.							
11. Quickly review the Five Rights once again while checking the epinephrine auto-injector to be sure that it is:							
a. For the right child							
b. The right medication							
c. The right dose (an approximation of the child's weight may be needed)							
d. Being given at the right time and							
e. Being given by the right route.							
12. Epi-Pen® Administration Remove the EpiPen auto-injector from the container.							
Remove the blue safety guard from the pen.							
Firmly press the orange tip against the student's mid-outer thigh until click is felt, and then hold for 3 (three) seconds.							

a. The pen can be given through clothing in most circumstances.							
Remove and massage the area for 10 seconds.							
Replace the pen back into its container and place it into an appropriate sharps container or hand to emergency medical responders.							
13. Impax/Amneal/Lineage Epinephrine Injection. USP Auto- Injector Administration Remove the auto- injector from its container							
Remove the caps at both ends of the auto-injector							
Firmly press the red rounded tip down hard against the student's mid-outer thigh until the needle penetrates and hold for 10 (ten) seconds. a. The injector can be given through clothing in most circumstances.							
Remove from the thigh and massage the area for 10 seconds. a. Take precaution with the exposed needle while replacing the injector back into the container.							
Dispose of the injector into an appropriate sharps container or hand to emergency medical responders							
14. AUVI-Q® Administration Pull AUVI-Q® up from the outer case.							
Pull red safety guard down and off of AUVI-Q®. a. The red safety guard is made to fit tightly. Pull firmly to remove. b. To reduce the chance of an accidental injection, do not touch the black base of the							

<p>auto-injector, which is where the needle comes out.</p> <p>c. If an accidental injection happens, get medical help right away.</p>							
<p>Place black end of AUVI-Q® against the middle of the outer thigh, then push firmly until you hear a click and hiss sound, and hold in place for 2 (two) seconds. AUVI-Q may be given through clothing if necessary</p>							
<p>Place the used AUVI-Q® into an appropriate sharps container or hand to emergency medical responders</p>							
<p>15. SYMJEPI® Administration Pull cap off to expose needle. a. Handle the device carefully to avoid an accidental needle stick.</p>							
<p>Holding by the finger grips, slowly insert the needle into the thigh. a. The injection can be given through clothing.</p>							
<p>After needle is in the thigh, push the plunger all the way down until it clicks.</p>							
<p>Hold in place for 2 seconds.</p>							
<p>Remove needle and massage the area for 10 seconds. a. Note: The correct dose has been injected if the plunger has been pushed all the way down and the solution window is at least partially blocked.</p>							
<p>Upon removing the needle be cautious of exposed needle and take care to avoid an accidental needle stick.</p>							
<p>While holding the plunger end of the syringe with one hand, use your other hand to slide the safety guard up until it clicks to cover the needle.</p>							

Place SYMJEPI® auto-injector into an appropriate sharps container or hand to emergency medical responders							
16. Teva Generic Administration Remove the Teva auto-injector from the container							
Twist off the cap in the direction of the arrow (yellow for high dose, green for low dose auto-injectors).							
Grasp device in fist and point downwards. Pull off blue safety release.							
Firmly press the orange tip against the student's mid-outer thigh until click is felt, and then hold for 3 (three) seconds. The injection may be given through clothing.							
Remove and massage the area for 10 seconds.							
Replace the pen back into its container and place it into an appropriate sharps container or hand to emergency medical responders.							
17. If alone with the student and you have not already called 911 or emergency medical services and notified parents, do so now.							
18. Monitor student's pulse and respirations.							
19. Once the emergency medical responders arrive, inform them of the medication administered, including the type of medication, dose, and the time given.							
20. If not done already, dispose of the injector in an approved sharps disposal container or give to the emergency medical responders.							
21. Remove gloves and wash hands.							

22. Document medication administration in student's medication administration log.							
23. Follow up with the parent or guardian and healthcare provider, as needed.							

Plan for monitoring medication administration:

School Nurse Name: _____ **Phone Number:** _____